


CCMC: Prototyping & Accelerating Implementation of Advanced Space Weather Models & Forecasting Systems.



Masha Kuznetsova & CCMC team

Space Weather Workshop April 8-11, 2014

MODELS • DATA • TOOLS • SYSTEMS • SERVICES • INNOVATIVE SOLUTIONS



#### **Outline**



- CCMC Now: brief overview.
- What CCMC is doing to accelerate implementation of advanced models.
- Focus on geospace models and partnership with NOAA/SWPC.
- Summary & outlook




Established in 2000 as an essential element of the National Space Weather Program

Designed to be a long-term solution to the R2O problem.

CCMC is fast and flexible with customer driven priorities.

Goals & functions are evolving in response to lessons learned & emerging needs.



Partnership is a key: with international research community, model owners, space weather operational agencies, NASA engineers and mission operators.



Collaborative

Development

LWS

Deliverables

Recipient

Missions

Science

Support

### **CCMC/SWRC** Activities at a Glance



## **Science Support**

Runs on Request

Visualization

& Analysis

Models

Data

Innovative **Solutions** 

For Research,

Analysis & **Forecasting** 

Testing & **Validation**  **R20 Support** 

Real-Time Systems:

Simulations. Data Streams,

**SW Widgets** 

Services

for NASA

missions

Forecasting

Methods

Scoreboard

**Environment** Monitoring by Experts Interplanetary

Space

**Space Weather** Notifications,

& Reports

**Anomaly** 

**Analysis** 

**Prototyping** 

Knowledgebase

Community **Projects** Leadership

Access & I/O Interpolation

Libraries

*Flexible* Dissemination

Database of SW Events

**Standards** 

**Database** of Run Results



#### **CCMC Staff**
















S. Bakshi

Anna Chulaki

(Director)


(Deputy)

P. Macneice

M. Mendoza







L. Rastaetter



**Space Weather Research Center** 

Ja Soon

Shim



M. Swindell



Chiu Wiegand



Kiran Patel



A. Taktakishvili



Yihua Zhen



Alex Gloce Leila Mays



A. Pulkkinen (Lead) Collado-Vega



## CCMC/SWRC Lab at NASA/GSFC Bld. 21, Rm 261






and a demo





## **Expanding Collection of Models**






> 60, from Sun to Earth and beyond




## **Expanding Collection of Models**





> 40, from L1 to Geospace



#### Magnetosphere Models Highlights:





#### Global Magnetosphere (physics-based)

- Space Weather Modeling Framework (SWMF) (Gombosi et al, U. Mich)
  - SWMF with RCM (v20110131) (selected by NOAA/SWPC)
  - SWMF with RCM+RBE (Radiation Belt Environment)
  - SWMF with CRCM (Comprehensive Ring Current Model)
- LFM Magnetosphere (Lyon et al, Dartmouth/NCAR/JHU-APL/CISM)
  - LFM-MIX
  - CMIT: fully coupled LFM-MIX & TIEGCM, version 2.2.0
- OpenGGCM v4.0 (Raeder, UNH)

#### Inner Magnetosphere (physics-based)

- RCM (Sazykin et al, Rice. U)
- Fok Models (M-C. Fok et al, GSFC)
  - Radiation Belt Environment (RBE)
  - Comprehensive Ring Current Model (CRCM)
  - CIMI: CRCM + RBE
- Plasmasphere model (Pierrard et al, IASB, Belgium)
- **VERB** radiation belt model (*Y. Shrits*) (test mode)



## CCMC Signature Services (Simulations)





#### Runs-on-Request Service

Web-base interactive system to serve advanced models to the international research community

- User-configurable input parameters and settings.
- Comprehensive on-line visualization & downloads

A tool for designing of sw products (widgets).

# Event-Triggered Simulations (real-time & historic)

#### FastTrack Request Option

- Input files are automatically downloaded.
- Optimal setting for real-time simulations.

Examples: Enlil Cone Model and/or SWMF simulations triggered when earth-directed CME is detected.

#### A 1-Click tool for forecasting and real-time validation

Continuous Real-Time Simulations

- Real-time data drivers.
- Data flow monitoring and controlling systems.


Example: SWMF (without RCM) since 2002.

Provide feeds to real-time widgets & displays.




# CCMC Signature Services: iSWA Integrated Space Weather Analysis System (Innovative Dissemination)






- Custom displays composed interactively by user from catalog of > 350 different widgets (cygnets), > 270 feeds from CCMC models.
- Cygnets are designed in collaboration with users of sw products & CCMC partners: NOAA/SWPC, AFWA, NASA mission operators.
- Combines simulations results & observational data (real-time & archives).


A multi-purpose tool for real-time space environment monitoring, event analysis, validation, education & forecaster training, collaboration.




## iSWA Cygnets: PBMOD (Ionosphere Scintillations)










J. Retterer, CCMC: J-S. Shim




## Comprehensive Inner Magnetosphere-Ionosphere (CIMI) Model: Real-Time Fluxes at VAP Orbit




#### Electrons at RBSP-A



#### 100 keV protons at RBSP (L-time)






#### iSWA Cygnets: Radiation Belt Environment (RBE) Model



RBE electron flux (> 0.6 MeV) at GEO W75

GOES electron flux (> 0.8 MeV)









### iSWA Cygnets: Realtime Observational Data




#### JRO Electron Density



#### Van Allen Electron Flux







### Global Magnetosphere Models Space Weather Widget Examples




Auroral boundaries and polar cap absoption




Joule heating integrated over polar cap




Magnetopause position



Ground magnetic perturbations, dB/dt. Primary interest of NOAA/SWPC (key to GICs).





#### Geospace Model Validation Efforts



Build upon GEM 2008 Modeling Challenge (community-wide effort).

Select physical parameter of interest to users: ΔB, dB/dt, Regional-K (critical role of NOAA/SWPC).


Agree on metrics format (threshold-based metrics parameters and skill score formula) (critical role of NOAA/SWPC).

Develop (and test) a post processing tool that derives ground magnetic perturbations from global magnetosphere model outputs from all current systems (*L. Rastaetter*, CCMC). Can be applied to all models.

Develop interactive on-line validation system, that

- allow access to simulation results archive,
- automatically calculates skill scores for selected events and stations,
- trace model performance over time.

Make validation process a seamless part of model development circle. Real-time prototyping and validation.




### Validation System for 6 Events



#### Groundbased magnetic perturbations dB/dt and Regional-K study results

Event	Magnetic perturbations on the ground dB/dt	Magnetic perturbations on the ground Delta B	Regional-K
2006/12/14 (doy 348) 12:00 UT - 12/16 00:00 UT	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
2001/08/31 (doy 243) 00:00 UT - 09/01 00:00 UT	ABK FRD FRN FUR IQA MEA NEW OTT PBQ WNG YKC	ABK FRD FRN FUR IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
2005/08/31 (doy 243) 10:00 UT - 09/01 12:00 UT	ABK FRD FRN FUR HRN MEA NEW OTT PBQ WNG YKC	ABK FRD FRN FUR HRN MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
2003/10/29 (doy 302) 06:00 UT - 10/30 06:00 UT	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
2010/04/05 (doy 095) 00:00 UT - 2010/04/06 00:00	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK NEW OTT SNK WNG YKC
2011/08/05 (doy 217) 09:00 UT - 2011/08/05 09:00	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK NEW OTT SNK WNG YKC

You can also download Delta B, dB/dt and K-index timeseries files.




### Validation System for 6 Events



#### Groundbased magnetic perturbations dB/dt and Regional-K study results

Event	Magnetic perturbations on the ground dB/dt	Magnetic perturbations on the ground Delta B	Regional-K
2006/12/14 (doy 348) 12:00 UT - 12/16 00:00 UT	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBO WNG YKC	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
B_North from observatory	file: new_OBS_20031029.txt  Observation: — new  Model runs: — 9a_SWMF — 4_OPENGGCM — 2_LFM-MIX	ABK FRD FRN FUR IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
ENorth -500		ABK FRD FRN FUR HRN MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
-1500 -2000 8:00 12:00 16:00	20:00 0:00 4:00	ABK FRD FRN FUR HRN IQA MEA NEW OTT PBQ WNG YKC	ABK NEW OTT PBQ WNG YKC
OO:OO UT - 2010/04/06 OO:OO	HRN IQA NEW OTT SNK WNG YKC	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK NEW OTT SNK WNG YKC
2011/08/05 (doy 217) 09:00 UT - 2011/08/05 09:00	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK FRD FRN FUR HRN IQA NEW OTT SNK WNG YKC	ABK NEW OTT SNK WNG YKC

You can also download Delta B, dB/dt and K-index timeseries files.



#### Run on Requests ΔB at 130 magnetometer positions for all magnetospere models

#### Paul Tenfiord 033114 4

Title/Introduction:

Key Word: sharp turn in B v

Model Type: GM

Model: LFM version LTR-2 2 0

Inflow Boundary Conditions: Start Time: 2000/01/01 00:00 End Time: 2000/01/01 03:04 Dipole Tilt at Start in X-Z Plane: 0.00 deg. Dipole Tilt in Y-Z GSE Plane: 0.00 deg. Dipole Update With Time: no Ionospheric Conductance: auroral

Radio Flux 10.7 cm: 150

Coordinate System for the Output: SM Initial Solar Wind (SW) Parameters in SM Coordinates:

SW Density: 8.00000 n/cc

SW Temperature [Kelvin]: 13335.20000 Kelvin X Component of SW Velocity: -400.00000 Y Component of SW Velocity: 0.00000 km/sec Z Component of SW Velocity: 0.00000 km/sec IMF Bx: 0.00000 nT

IMF By: 0.00000 nT IMF Bz: 0.00000 nT IMF |B|: 0.00000 nT IMF Clock Angle: o deg.

- View solar wind input data
- · List solar wind input data in ASCII format (see format desc
- View Magnetosphere
- Create Timeseries in Magnetosphere
- · View Ionosphere

#### View pre-computed timeseries data:

- · Northern hemisphere polar cap flux and area
- Southern hemisphere polar cap flux and area
- Magnetopause standoff and closest approach within 30 deg. of Sun-Earth line (local noon)
- Polar cap boundary at 24 magnetic local times in northern hemisphere
- Polar cap boundary at 24 magnetic local times in southern hemisphere
- Ionospheric dissipation
- · View Ouick look graphics for the run


View Magnetic perturbation calculated for magnetometer stations:



B\_NorthGeomag B\_EastGeomaa FUR\_magnetometer.dat [nŤ [nT] 10,00 5.0 10.0 5.00 Model: LFM 0.0 e-dimensional model output. -5.0anges. 0.00 ing image will be displayed at this location -10.0-10.0tered a programming error. Please report -5.00-20.00:00 1:00 2:00 3:00 Second: 0 Second: 59,99998 2000/01/01 03:41 1999/12/31 1:00:00 time Model at CCMC: LFM Log scale (apply to all quantities > 0 in plot) Lock plot data range: Min.: 0 Max.: 1 Image magnification: 1 Line style: no line \* Plot symbols: diamonds Symbol size: 0.2 Reset Form will reset changes to the defaults specified by the previous run of this script. Update Plot will update (generate) the plot with the chosen time and plot parameters above or will print the entire file to screen. Have data printed to text file. The file format is slightly different from the original file visualized and can include computed quantities if offered and selected for plotting (i.e. quantities beyond basic MHD quantities N, P, V, B), such as B, V, J, E, Jpar.


> Runs-on-Request: Contact CCMC Staff Visualization: Dr. Lutz Rastätter

Last script update: August 18, 2011.



#### iSWA: SWMF (BATSRUS + RCM) v. 20110131 (U. Mich)






DST, Ground peturbations (delta-B, dB/dt) timelines - coming soon.



### Sensitivity Analysis to Station Location








### **Ensemble of Virtual Magnetometers**








## **Ensemble of Virtual Magnetometers**









### **Ensemble of Virtual Magnetometers**






#### **Ensemble Displays (Event Simulations)**




**FastTrack** option with automated skill score calculation to trace model improvement over time – *coming soon*.




## Pushing Models to the Limits: Modeling Extreme SW Event




- Artificially constructed solar wind driving conditions (consistent with available information on Carrington storm).
- Specially adapted version of the SWMF:
  - inner boundary at 1.25 Re vs 2.5 Re,
  - resolution at inner boundary 1/16 Re vs 1/4 Re.
- The inner boundary condition becomes extremely important during periods of strong driving.





Working with code developers to address the issue

C. Ngwira, A. Glocer, G. Toth



### Partnership with NASA Users



## Linking CCMC space environment modeling systems with models calculating impacts on systems in space


JSC/SRAG, D. Fry et al, Space radiation analysis

JPL, H. Garret et al., Surface charging modeling

MSFC, J. Minow et al., Internal charging

GSFC, K. Label et al., SEU modeling

GSFC, L. Newman, Satellite drag & conjunction assessment




#### Summary and Outlook



- CCMC is designed for rapid implementation and prototyping of advanced capabilities (including LWS deliverables).
- On-going activities in support of geospace model transition to NOAA/SWPC:
  - Continuous real-time SWMF runs with results dissemination through iSWA displays and Web services.
  - Event triggered simulations for SWMF and other global magnetosphere models (LFM, OpenGGCM).
  - Automated dB/dt and skill score calculations.
  - Push models to the limits. Sensitivity analysis.
- Operation agencies are invited to take advantage of iSWA.
  - ✓ Test-drive before investing.
  - ✓ Opportunity for collaborative tool development.
- Participation of users of space weather products in setting validation projects is a key.
- Much more is possible if we continue working together and expand collaboration.






## **Supplementary Material**






## An extensible software system for NASA that provides an unprecedented capability for:


- viewing space environment conditions for specific missions/orbits
- providing automated SW alerts
- assimilating and displaying spacecraft anomaly information
- displaying spacecraft/mission data







## Collaborative Development with Model Owners



All drivers are converted to a common format.

The tool is called as a KAMELEON subroutine to provide values on the grid;

call kameleon (model, time, mlts mlats, variables, values\_output)

J-S. Shim M. Swindell

IT Models: CTIPe/TIE-GCM/GITM